

On parametric phenomena in correspondence theory

Willem Conradie

*School of Mathematics, University of the Witwatersrand, Johannesburg,
National Institute for Theoretical and Computational Sciences (NITheCS), South Africa*

Abstract

I survey a number of recent results which study the syntactic shape of modal formulas' first-order frame correspondents on different relation semantics and establish systematic links between the correspondents of given modal formula across these different semantic environments. It has been observed that, for instance, in some cases, correspondents remains syntactically unchanged under such semantic shifts; in other cases, correspondents across different semantics can be recovered as parametric instances of the same pattern, with the parameter capturing the embedding between semantic contexts. I discuss some open problems and further directions.

Keywords: Correspondence theory, modal logic, many-valued modal logic, intuitionistic modal logic, non-distributive logics.

Unified correspondence. In recent years, the research programme of unified correspondence (see, e.g., [9] [14]) has developed Sahlqvist correspondence and canonicity results for large families of non-classical logics. In [10] we formulate a general definition of the classes of Sahlqvist and inductive formulas which applies uniformly to any logic algebraically captured by a class of lattice expansions (LE-logics). This definition is based solely on the order-theoretic properties of the algebraic interpretation of the connectives and, when projected onto specific LE-logics, captures or generalises many existing definitions in the literature. A further feature of unified correspondence, is a division of labour which separates the computation of a first-order correspondent for a modal axiom into two phases: a correspondent is first computed and expressed as a conjunction of quasi-inequalities in an extended modal language interpreted on perfect lattice expansions. This is independent of any particular choice of relational semantics for the logic. The second phase of the computation translates this algebraic correspondent by applying the standard translation associated with a particular choice of relational semantics linked with the algebraic semantics via a suitable duality.

Parametric correspondence. This and other recent work in correspondence theory, has focused mostly on the (modal) propositional side of the correspondence theoretic equation, i.e., on identifying classes of (modal) propositional formulas for which first-order correspondents exists. However, beyond the fact of their existence, the properties of the first-order correspondents themselves have received little attention. This situation has changed recently, with a number of results studying the syntactic shape of first-order correspondents on particular relation semantics and, more specif-

ically, comparing the shape of the first-order correspondent of given modal formula across different relational semantics. It has been observed, for instance, that, in some cases, correspondents remains syntactically unchanged under such semantic shifts; in other cases, correspondents across different semantics can be recovered as parametric instances of the same pattern, with the parameter capturing the embedding between semantic contexts. This talk will survey a range of phenomena and results of this type, some of which I discuss below.

Many-valued modal logic. The first result of this type was proven in the context of the correspondence theory of many-valued modal logic (see, e.g., [12,3]). Specifically, considering many-valued modal logics with complete Heyting algebras as truth value spaces, [4] shows that the first-order correspondent of any modal Sahlqvist formula over many-valued Kripke frames is syntactically identical to its first-order correspondent on (crisp) Kripke frames. Of course, the many-valued correspondent should be read as a formula of many-valued predicate logic and therefore its semantic meaning generalizes that of the classical correspondent. Apart from purely theoretical interest, this result has great practical utility as it means that the correspondents of well-known modal axioms can be directly transferred to the many-valued versions of the logics they axiomatize.

Non-distributive modal logics with polarity-based semantics. The next important instance of parametric correspondence which I would like to discuss comes from the study of non-distributive modal logics [11]. These logics can be given semantics based on polarities, or formal contexts, in the sense of formal concept analysis (FCA) [13] and given an epistemic interpretation [8,7] where formulas denote formal categories. In [8,7] it is observed that, over this semantics, the classical S5 axioms of epistemic logic have first-order correspondents which are, in a sense, dual to their well-known correspondents on Kripke frames. Moreover, it is argued that the epistemic meanings of these axioms are preserved in a modified form in this new context. This observation is extensively developed in [6] where the first-order correspondents of modal reduction principles [18] are captured as inclusions of relational compositions, both in the setting of Kripke frames and in that of polarity based frames. It is established that one may pass from the Kripke frame correspondent, thus expressed, to the correspondent on polarity-based frames, roughly speaking, by reversing the inclusion and parameterising the relational compositions with the incidence relation of the polarity. Mathematically, this is underpinned by the embedding of the class of Kripke frames into the class of enriched polarities.

Non-distributive modal logics with graph-based semantics. Alternatively, non-distributive logics may be given graph-based semantics [11,5] based on Ploščica's [15] representation of general lattices. Here too, the correspondents of modal reduction principles may be obtained by appropriately transforming their correspondents on Kripke frames.

Intuitionistic modal logic. In the case of intuitionistic modal logic with semantics based on [19], the correspondents of inductive (and hence also Sahlqvist) formulas can be transparently obtained from the classical correspondents. This is mainly due to the strong interaction conditions between the pre-order of the intuitionistic Kripke

frame and the additional accessibility relation used to interpret modalities. In the case of Fisher-Servi's semantics for intuitionistic modal logic [16,17], the interaction conditions between the relations are weaker, and the relationship between correspondence in the classical case and in this setting is somewhat more involved.

Relativized correspondence. When imposing restrictions on the class of Kripke frames considered, more modal formulas become first order definable, in a phenomenon known as relativized correspondence. For instance, all modal reduction principles are first-order definable over transitive frames [18], while all modal formulas have first-order correspondents over S5 frames [2] and over Euclidean frames [1]. Relativized correspondence has been little studied for modal logics on non-classical propositional base or for many-valued modal logics, but some initial results indicates that an interesting landscape waits to be discovered here. I will conclude by sketching some of these results and directions.

References

- [1] Balbiani, P., D. Georgiev and T. Tinchev, *Modal correspondence theory in the class of all euclidean frames*, Journal of Logic and Computation **28** (2018), pp. 119–131.
- [2] Balbiani, P. and T. Tinchev, *Definability over the class of all partitions*, Journal of Logic and Computation **16** (2006), pp. 541–557.
- [3] Bou, F., F. Esteva, L. Godo and R. O. Rodríguez, *On the minimum many-valued modal logic over a finite residuated lattice*, Journal of Logic and computation **21** (2011), pp. 739–790.
- [4] Britz, C., “Correspondence theory in many-valued modal logics,” Master's thesis, University of Johannesburg, South Africa (2016).
- [5] Conradie, W., A. Craig, A. Palmigiano and N. Wijnberg, *Modelling informational entropy*, in: *Proc. WoLLIC 2019*, Lecture Notes in Computer Science **11541** (2019), pp. 140–160.
- [6] Conradie, W., A. De Domenico, K. Manoorkar, A. Palmigiano, M. Panettiere, D. P. Prieto and A. Tzimoulis, *Modal reduction principles across relational semantics*, arXiv preprint arXiv:2202.00899 (2022).
- [7] Conradie, W., S. Frittella, A. Palmigiano, M. Piazzai, A. Tzimoulis and N. Wijnberg, *Toward an epistemic-logical theory of categorization*, in: *16th conference on Theoretical Aspects of Rationality and Knowledge (TARK 2017)*, Electronic Proceedings in Theoretical Computer Science **251**, 2017, pp. 170–189.
- [8] Conradie, W., S. Frittella, A. Palmigiano, M. Piazzai, A. Tzimoulis and N. M. Wijnberg, *Categories: how I learned to stop worrying and love two sorts*, in: *Proc. WoLLIC 2016*, Lecture Notes in Computer Science **9803**, Springer, 2016, pp. 145–164.
- [9] Conradie, W., S. Ghilardi and A. Palmigiano, *Unified Correspondence*, in: A. Baltag and S. Smets, editors, *Johan van Benthem on Logic and Information Dynamics*, Outstanding Contributions to Logic **5**, Springer International Publishing, 2014 pp. 933–975.
- [10] Conradie, W. and A. Palmigiano, *Algorithmic correspondence and canonicity for non-distributive logics*, Annals of Pure and Applied Logic **170** (2019), pp. 923–974.
- [11] Conradie, W., A. Palmigiano, C. Robinson and N. Wijnberg, *Non-distributive logics: from semantics to meaning*, in: A. Rezus, editor, *Contemporary Logic and Computing*, Landscapes in Logic **1**, College Publications, 2020 pp. 38–86.
URL arXivpreprintarXiv:2002.04257
- [12] Fitting, M., *Many-valued modal logics*, Fundam. Inform. **15** (1991), pp. 235–254.
- [13] Ganter, B. and R. Wille, “Formal concept analysis: mathematical foundations,” Springer Science & Business Media, 2012.
- [14] Greco, G., M. Ma, A. Palmigiano, A. Tzimoulis and Z. Zhao, *Unified correspondence as a proof-theoretic tool*, Journal of Logic and Computation **28** (2018), pp. 1367–1442.
- [15] Ploščica, M., *A natural representation of bounded lattices*, Tatra Mountains Mathematical Publications **4** (1994).

- [16] Servi, G. F., *Semantics for a class of intuitionistic modal calculi*, in: *Italian studies in the philosophy of science*, Springer, 1980 pp. 59–72.
- [17] Servi, G. F., *Axiomatizations for some intuitionistic modal logics*, *Rend. Sem. Mat. Univers. Politecn. Torino* **42** (1984), pp. 179–194.
- [18] Van Benthem, J., *Modal reduction principles*, *The Journal of Symbolic Logic* **41** (1976), pp. 301–312.
- [19] Wolter, F. and M. Zakharyashev, *Intuitionistic modal logic*, in: *Logic and foundations of mathematics*, Springer, 1999 pp. 227–238.